



This is a repository copy of *Contextual and individual factors associated with dental services utilisation by Brazilian adults: A multilevel analysis*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/128200/>

Version: Published Version

Article:

Herkath, F.J. orcid.org/0000-0003-4439-0189, Vettore, M.V. orcid.org/0000-0001-6299-4432 and Werneck, G.L. (2018) Contextual and individual factors associated with dental services utilisation by Brazilian adults: A multilevel analysis. PLoS One, 13 (2). e0192771.

<https://doi.org/10.1371/journal.pone.0192771>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

RESEARCH ARTICLE


Contextual and individual factors associated with dental services utilisation by Brazilian adults: A multilevel analysis

Fernando José Herkrath^{1,2,3☯*}, Mario Vianna Vettore^{4☯}, Guilherme Loureiro Werneck^{2,5☯}

1 Escola Superior de Ciências da Saúde, Universidade do Estado do Amazonas, Manaus, Amazonas, Brazil, **2** Instituto de Medicina Social, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil, **3** Instituto Leônidas e Maria Deane, Fundação Oswaldo Cruz, Manaus, Amazonas, Brazil, **4** Academic Unit of Oral Health, Dentistry and Society, School of Clinical Dentistry, University of Sheffield, Sheffield, United Kingdom, **5** Instituto de Estudos em Saúde Coletiva, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

☯ These authors contributed equally to this work.

* fernandoherkrath@gmail.com



Abstract

Background

Inequalities in the utilisation of dental services in Brazil are remarkable. The aim of this study was to evaluate the association of contextual and individual factors with the utilisation of dental services by Brazilian adults using the Andersen's behavioural model.

Methods

Individual-level data from 27,017 adults residents in the State capitals who were interviewed in the 2013 Brazilian National Health Survey were pooled with contextual city-level data. The outcomes were non-utilisation of dental services and last dental visit over 12 months ago. Individual predisposing variables were age, sex, race/skin colour, schooling and social network. Individual enabling variables included income, health insurance and registration in primary health care. Individual need variables were self-perceived dental health and self-reported missing teeth. Multilevel logistic regression models were performed to estimate odds ratio (OR) and 95% confidence intervals (95% CIs) of the association of contextual and individual predisposing, enabling and need-related variables with dental services outcomes.

Results

Predisposing (OR = 0.89; 95% CI 0.81–0.97) and enabling (OR = 0.90; 95% CI 0.85–0.96) contextual factors were associated with non-utilisation of dental services. Individual predisposing (sex, race/skin colour, schooling), enabling (income, health insurance) and need (self-perceived oral health, missing teeth) were associated with non-utilisation of dental services and last dental visit over 12 months ago. The latter was also associated with other individual predisposing (age, social network) and need (eating difficulties due to oral problems) characteristics.

OPEN ACCESS

Citation: Herkrath FJ, Vettore MV, Werneck GL (2018) Contextual and individual factors associated with dental services utilisation by Brazilian adults: A multilevel analysis. PLoS ONE 13(2): e0192771. <https://doi.org/10.1371/journal.pone.0192771>

Editor: Peter M. Milgrom, University of Washington, UNITED STATES

Received: October 5, 2017

Accepted: January 30, 2018

Published: February 8, 2018

Copyright: © 2018 Herkrath et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: National Health Survey data is in the public domain and made available from the Instituto Brasileiro de Geografia e Estatística (IBGE) database (ftp://ftp.ibge.gov.br/PNS/2013/microdados/pns_2013_microdados_2017_03_23.zip).

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: SUS, Brazilian Unified Health System; FHT, family health teams; NHS, national health survey; IBGE, Brazilian Institute of Geography and Statistics; PSU, primary sampling unity; VPC, variance partition coefficient.

Conclusions

Individual and contextual determinants influenced dental services utilisation in Brazilian adults. These factors should be on the policy agenda and considered in the organisation of health services aiming to reduce oral health inequalities related to access and utilisation of dental services.

Introduction

Public oral health care in Brazil has been historically restricted to a limited range of dental procedures offered in the large urban centres. In this context, curative dental procedures have been predominating over preventive dental services and access to oral health care has been restricted to few population groups [1]. However, over the recent years, the organisation of public dental services has been improved with the implementation of the National Oral Health Policy. The core principles of the policy focus on the reorganisation and expansion of primary oral health care within the Brazilian Unified Health Care System (SUS). In addition, the policy framework proposes the increase in the provision of universal dental care through combining actions at individual and collective levels encompassing health promotion, prevention, diagnosis, treatment and rehabilitation [2].

Since 1998 primary care in the SUS has been reorganised around a priority strategy which provides community-based primary health care through Family Health Teams (FHT), that are responsible for the health of the population at a defined geographical area [3, 4]. The gradual incorporation of oral health professionals in the teams occurred since 2000. The implementation of the National Oral Health Policy was accompanied by a significant increase in the federal public funding, resulting in the expansion of access and enhancing comprehensiveness of health care [1, 5–7]. This propitiated the expansion of oral health teams in primary care and the implementation of dental specialties centers for secondary care, offering procedures that were previously exclusive to private services [1, 5].

However, despite the increase in access, large inequalities in dental health care utilisation and in the oral health status of the Brazilian population remain noticeable. The most vulnerable groups, such as rural, elderly, deprived and less educated populations, face more barriers in using health services and present have worse oral health conditions. Furthermore, epidemiological surveys also revealed marked oral health inequalities between the regions of the country [6–16].

Utilisation of health services results from a complex interaction between users, availability and access of services and is influenced by several factors, including individual characteristics (e.g. perception of health status and health needs), contextual factors (e.g. social inequalities) and organisation of health services (e.g. distribution of health care units) [17–19]. Previous studies on the associated factors highlight that dental services utilisation results from the interaction between biological, sociocultural, family and community determinants, as well as from the characteristics of health systems [20, 21]. The literature also suggests that reducing social inequalities plays an important role in mitigating barriers to health services utilisation [7, 22–24].

Dental care utilisation and frequency of dental services utilisation by Brazilian adults have been associated with high schooling [7, 8, 10, 12, 22, 25–32] and better-off socioeconomic status [8, 12, 26–29, 32]. Females [7, 8, 26–30], younger people [7, 8, 22, 26, 28, 33], white race/skin colour [7, 34] and having a health insurance [7, 27, 30] have also been associated with a

greater utilisation of dental care. Regular users have less dental treatment needs and better oral health conditions [12, 22, 25, 26, 29, 31], and better general and oral health self-perception [26, 28, 29, 31, 32]. Contextual factors, such as living in urban areas [8, 33] and in locations covered by the FHT in primary care, have been linked with an increased likelihood of health services utilisation [7].

Similar findings regarding inequities in the oral health conditions and utilisation of services have been reported in other countries. Populations and countries with greater social inequalities present higher burden of oral diseases and lower use of dental services [20, 23, 24, 35–42]. Investigating the associated factors of utilisation of dental services is relevant for planning realistic interventions to improve access and quality of health care, reorganisation of health services, and ultimately to improve the use of available resources [21, 43]. The identification of the determinants of non-utilisation dental services as a way of assessing barriers to access to dental services can contribute to the improvement of equity in oral health care, but recent data from Brazil have not yet been extensively explored for this purpose. Thus, the aim of the study was to evaluate the association of contextual and individual factors with the utilisation of dental services by Brazilian adults living in the State capitals.

Methods

Study design

This cross-sectional study used secondary data from the Brazilian National Health Survey (NHS), a nationwide home-based survey conducted in 2013 by the Ministry of Health in partnership with the Brazilian Institute of Geography and Statistics (IBGE). The purpose of NHS was to evaluate the health status and lifestyles of the Brazilian population, as well as health care, including utilisation and access to health services, continuity of care and financing [44]. All data are fully available without restriction available through IBGE homepage (<https://ww2.ibge.gov.br/english>) and database (ftp://ftp.ibge.gov.br/PNS/2013/microdados/pns_2013_microdados_2017_03_23.zip).

Study population

The NHS sample was representative for Brazil, geopolitical macro-regions, States, metropolitan regions and States capitals. Census tracts or a set of these with at least 60 households was defined as the primary sampling unity (PSU). Sampling was performed in three stages. In the first stage, census tracts were selected with probability proportional to size (given by the number of households in each unit) in each stratum. A simple random sampling of households was selected in the second stage in each PSU selected in the first stage. The third stage was a simple random sampling of one adult aged 18 years or older among all adults living in the household to answer a comprehensive questionnaire. The present study analysed data from 27,017 adult dwellers in the 27 State capitals.

Study variables

The two outcome variables related to utilisation of dental services were based on the following question: “When was your last dental appointment?”, with the following response options: 1. In the last 12 months; 2. From one year to less than two years; 3. From two years to less than three years; 4. Three years or more; 5. Never had a dental appointment. The categories were grouped in order to derive two outcomes: (i) non-utilisation of dental care, considering the option “never had a dental appointment” as outcome (response option 5 vs 1–4); and (ii) last dental visit over 12 months ago, assessed only among those who have already been to the

dentist (response options 2–4 vs 1, those who answered option “5” were excluded from this analysis).

The selection of independent variables was based on Andersen’s Behavioural Model of Health Services Utilisation [43]. Individual predisposing characteristics included age, sex, self-reported race/skin colour, according to the categories used in Brazil by the IBGE (white, black, yellow/Asiatic, brown/“pardo” and indigenous) [45], schooling (years of study completed with approval) and social network. Social network was assessed using the items proposed by Berkman and Syme [46], categorising individuals into four groups: I, being the more isolated category, and II, III or IV composing the other three categories of gradation in social relations. Individual monthly income, health insurance and registration in primary care FHT were selected as enabling factors. Perceived need included self-perception of oral health and self-reported difficulty to eat due to oral problems and evaluated need was assessed through self-reported number of missing teeth.

Contextual variables included city-level predisposing, enabling and need indicators. Predisposing contextual characteristics included information about social vulnerability for the year 2010, including data on income, education and life expectancy from the Human Development Index obtained from the Atlas of Human Development in Brazil, besides proportion of individuals living in extreme poverty (household *per capita* monthly income \leq R\$70.00) and proportion of individuals vulnerable to poverty (household *per capita* monthly income \leq R\$255.00) [47]. Enabling factors were *per capita* expenditure in primary health care, *per capita* expenditure in oral health programs, and coverage of oral health teams in primary health care obtained from health information systems for the year 2013 [48]. The contextual need variables were number of permanent decayed, missing and filled teeth (DMFT index), self-reported dental pain in the last six months, need for dentures and impact of oral conditions on daily activities (OIDP questionnaire) among adults who participated at the 2010 Brazilian oral health survey. This information were collected at individual level and then aggregated at the city level [49].

Data analysis

First, the independent variables were described according to the dental services outcomes using proportions and 95% confidence intervals (95% CI), considering the complex sampling design and the sample weights (svyset UPA_PNS [pw = v0029], strata(v0024) poststrata(v00293) postweight(v00292)).

Second, factor analysis was used considering that contextual variables at city-level ($n = 27$ State capitals) are correlated and represent underlying dimensions related to utilisation of dental services. The oblique rotation method was used considering the existence of a correlation between the contextual variables. The predicted scores of the factors were used in subsequent analyses.

Third, multilevel logistic regression analysis with robust variance was performed to estimate the odds ratios (OR) and 95% CIs of contextual and individual independent variables (fixed effects with random intercept) and the two outcomes: (i) non-utilisation of dental services and (ii) last dental visit over 12 months ago. Logistic regression models were used since they provide more valid estimates of the incidence density ratio when cross-sectional data is analysed [50]. Initially, associations between independent variables and the two outcomes were tested in bivariate analysis. Variables that presented with $p < 0.10$ in bivariate analysis were considered for the multivariate hierarchical multilevel analysis. Stepforward non-automatic selection of variables in four models was conducted to obtain a parsimonious model according to Andersen’s theoretical model. The first model included only the contextual

factors. The second and third models consisted of individual predisposing and enabling variables, respectively. The final model included individual need-related variables. Variable that remained statistically significant at 10% ($p < 0.10$) were retained in the analysis for adjustment in the next model. The percentage of the variance explained by the contextual level (variance partition coefficient) was calculated using the standard logistic distribution variance, $\pi^2/3 \approx 3.29$, as the level 1 variance [51]. All analyses were performed in Stata 14.0 (College Station, TX USA).

Ethical issues

This study was approved by the Ethics Committee in Research of the State University of Rio de Janeiro, protocol CAAE number 64756517.0.0000.5260.

Results

The prevalence of non-utilisation of dental health services and last dental visit over 12 months ago by Brazilian adults from the State capitals was 1.8% (95% CI 1.6%–2.1%) and 48.1% (95% CI 47.0%–49.1%), respectively. [Table 1](#) presents the individual characteristics according to dental services utilisation outcomes. Most participants were females, had white race/skin colour, had 9–11 years of schooling and individual family income between R\$ 501 and R\$ 1500.

Factor analysis evidenced four factors that explained 89% of the total variance. [Table 2](#) presents the items loadings after oblique rotation. Bartlett's test for sphericity was statistically significant ($p < 0.001$). The model adequacy analysis using the Kaiser-Meyer-Olkin statistic was 0.612. The contextual factors were named as following: 1. contextual predisposing; 2. contextual enabling; 3. contextual perceived need; 4. contextual evaluated need.

In the unadjusted analyses, the predisposing contextual factor was associated with non-utilisation of dental health services and last dental visit over 12 months ago. The enabling contextual factor was also associated with non-utilisation of dental health services. In addition, all individual predisposing, individual enabling (except enabling financing) and individual need variables were associated with both outcomes ([Table 3](#)).

In the multilevel analyses, null logistic models were fitted in order to verify the presence of contextual level effects. Null models showed statistical significant random effects for both outcomes, demonstrating there were significant between-city differences. The percentage variance explained by the contextual level was 8.9% for the non-utilisation of dental services and 3.6% for last dental visit over 12 months ago. These proportions are reduced to 2.6% and 1.5% with the inclusion of contextual variables. The average number of observations in each cluster on the second level was 1,000.6, ranging from 520 to 2,568.

The multilevel logistic regression models for the outcome non-utilisation of services are presented in [Table 4](#). Predisposing and enabling contextual variables were inversely associated with non-utilisation of dental services in model 1 and remained associated afterwards. Individual predisposing demographic and social variables in model 2, and individual enabling financing in model 3 were associated with non-utilisation of dental services. In the final model (model 4), adults living in the cities with greater levels of contextual predisposing (OR = 0.77; 95% CI 0.64–0.94) and enabling factors (OR = 0.84; 95% CI 0.75–0.94) had significantly lower odds of non-utilisation of dental services. Sex, race/skin colour, schooling, income, health insurance, perceived oral health and missing teeth remained associated with non-utilisation of dental services.

[Table 5](#) presents the multilevel logistic regression analysis of the relationship between independent variables and last dental visit over 12 months ago. Contextual predisposing variable was associated with the outcome only in model 1. Individual predisposing and individual

Table 1. Individual predisposing, enabling and need characteristics according to non-utilisation of dental services and last dental visit over 12 months ago.

Variable	Total	Non-utilisation of dental services		Last dental visit > 12 months	
	%	%	95% CI	%	95% CI
<i>Predisposing demographic</i>					
Age					
18–21	8.5	2.8	1.9–4.1	40.7	37.5–44.0
22–34	29.2	1.8	1.4–2.4	44.5	42.8–46.3
35–44	19.8	1.9	1.3–2.7	45.3	43.4–47.2
45–64	30.0	1.4	1.1–1.9	49.0	47.1–50.8
≥ 65	12.5	2.0	1.5–2.6	63.5	60.9–65.9
Sex					
Male	45.0	2.4	2.0–2.9	51.8	50.2–53.4
Female	55.0	1.4	1.1–1.7	45.0	43.7–46.3
<i>Predisposing social</i>					
Race/skin colour					
White	47.2	1.0	0.7–1.3	43.5	42.0–45.0
Black	10.0	2.8	2.1–3.8	54.9	52.0–57.8
Yellow	1.5	1.1	0.3–4.0	39.3	31.0–48.3
Brown	40.8	2.6	2.1–3.0	52.0	50.5–53.5
Indigenous	0.5	7.9	3.2–18.0	47.8	37.4–58.4
Years of schooling					
0–4	15.8	4.9	4.1–5.9	64.9	62.7–66.9
5–8	19.2	2.4	1.9–3.0	60.7	58.7–62.7
9–11	38.1	1.4	1.1–1.9	46.4	44.9–48.0
> 11	26.9	0.2	0.1–0.3	31.5	29.8–33.2
Social network					
I (Low)	30.5	2.9	2.4–3.5	55.5	53.7–57.2
II	34.7	1.7	1.3–2.1	48.6	47.0–50.3
III	26.6	1.3	0.9–1.8	42.3	40.1–44.2
IV (High)	8.2	0.4	0.2–0.6	36.6	33.9–39.4
<i>Enabling financing</i>					
Income (R\$)					
≤500	7.1	4.2	3.2–5.5	54.2	50.8–57.6
501 a 1500	43.6	2.3	1.9–2.9	54.5	53.1–55.8
1501 a 2500	12.1	0.8	0.5–1.2	45.3	42.8–47.9
>2500	20.1	0.1	0.1–0.4	32.9	30.9–35.0
Not available	17.1	2.3	1.7–3.0	49.1	46.8–51.4
Health insurance					
No	57.4	3.0	2.6–3.5	56.9	55.6–58.1
Yes	42.6	0.2	0.1–0.3	36.2	34.8–37.7
<i>Enabling organisation</i>					
Registered in FHT					
No	38.3	1.9	1.5–2.4	51.2	49.7–52.7
Yes	50.0	1.7	1.4–2.1	46.0	44.5–47.6
Unknown	11.7	2.2	1.6–3.1	46.6	43.9–49.3
<i>Need perceived</i>					
Perceived dental health					

(Continued)

Table 1. (Continued)

Variable	Total	Non-utilisation of dental services		Last dental visit > 12 months	
	%	%	95% CI	%	95% CI
Very good / good	70.6	1.5	1.2–1.8	43.2	42.1–44.5
Regular	24.6	2.3	1.8–2.8	58.4	56.6–60.3
Poor / very poor	4.8	5.0	3.6–6.9	65.9	61.4–70.1
Eating difficulties due to oral problems					
No	92.1	1.6	1.4–1.9	47.2	46.1–48.3
Yes	7.9	4.1	3.1–5.5	58.3	55.3–61.1
Need evaluated					
Missing teeth					
No	33.2	2.1	1.7–2.7	43.0	41.3–44.8
One or more	58.6	1.5	1.3–1.9	46.4	45.1–47.7
All	8.2	2.8	2.0–3.8	80.6	78.0–82.9

<https://doi.org/10.1371/journal.pone.0192771.t001>

enabling variables were inserted in models 2 and 3, respectively. In the final model (model 4), the odds of visiting a dentist over 12 months ago was significantly higher for older adults, male sex, brown race/skin colour, low schooling, low social networks, low income, absence of health insurance, poor perceived dental needs and higher number of missing teeth (Table 5).

Discussion

This study showed the influence of contextual characteristics of social vulnerability and health services organisation in the non-utilisation of dental health services in Brazilian adults. However, these associations were not identified for last dental visit over 12 months ago. Individual predisposing, enabling and the need characteristics were associated with both outcomes of dental services utilisation. Sex, race/skin colour, schooling, income, health insurance, perceived dental health and tooth loss were associated with both outcomes. Age, social network

Table 2. Rotated component matrix presenting factor loadings for the contextual variables included in the factor analysis at city-level.

Variables	Component				Unexplained
	Predisposing	Enabling	Perceived need	Evaluated need	
HDI—income	0.947				0.088
HDI—life expectancy	0.897				0.125
HDI—education	0.807				0.281
Percentage of extremely poor	-0.972				0.067
Percentage of vulnerable to poverty	-0.978				0.047
Per capita expenditure in primary care		0.944			0.093
Per capita expenditure in oral programs		0.956			0.078
Family oral health teams coverage		0.976			0.054
Oral Impacts on Daily Performances			0.914		0.122
Dental pain			0.940		0.116
DFMT index				0.876	0.175
Need for denture				0.707	0.362

HDI: Human Development Index

Rotation method was oblique with Kaiser normalisation.

<https://doi.org/10.1371/journal.pone.0192771.t002>

Table 3. Bivariate analyses between independent variables and non-utilisation of dental services and last dental visit over 12 months ago, Brazil, State capitals, 2013.

Variables	Non-utilisation			Last dental visit > 12 months		
	OR	95% CI	p-value	OR	95% CI	p-value
Contextual (city-level)						
Predisposing (greater scores indicate better conditions)	0.63	0.52–0.76	<0.001	0.87	0.77–0.99	0.029
Enabling (greater scores indicate better conditions)	0.82	0.71–0.96	0.011	0.94	0.77–1.14	0.536
Perceived need (greater scores indicate worse perceived oral health)	0.94	0.80–1.11	0.469	0.98	0.93–1.03	0.408
Evaluated need (greater scores indicate worse oral health)	0.86	0.67–1.12	0.266	1.09	0.97–1.23	0.148
Individual level						
Predisposing demographic						
Age (ref.: 18–21)						
22–34	0.64	0.50–0.84	0.001	1.15	1.04–1.26	0.007
35–44	0.51	0.35–0.74	<0.001	1.22	1.09–1.36	<0.001
45–64	0.64	0.48–0.87	0.004	1.53	1.36–1.73	<0.001
65+	1.07	0.69–1.67	0.759	2.91	2.49–3.39	<0.001
Sex (ref.: female)						
Male	1.67	1.46–1.91	<0.001	1.30	1.24–1.37	<0.001
Predisposing social						
Race/skin colour (ref.: white)						
Black	2.71	2.23–3.30	<0.001	1.44	1.31–1.58	<0.001
Yellow	1.01	0.36–2.84	0.983	0.98	0.73–1.30	0.867
Brown	2.05	1.74–2.41	<0.001	1.38	1.28–1.49	<0.001
Indigenous	5.04	2.18–11.68	<0.001	1.10	0.78–1.56	0.596
Years of schooling (ref.: > 11)						
0–4	24.30	13.67–43.20	<0.001	4.79	4.30–5.33	<0.001
5–8	12.04	7.01–20.67	<0.001	3.35	3.02–3.72	<0.001
9–11	5.11	2.92–8.95	<0.001	1.78	1.62–1.96	<0.001
Social network (ref.: IV, High)						
I (Low)	4.20	2.20–8.01	<0.001	2.34	2.06–2.65	<0.001
II	2.53	1.36–4.70	0.003	1.64	1.47–1.84	<0.001
III	2.06	1.13–3.75	0.019	1.27	1.12–1.43	<0.001
Enabling financing						
Income (ref.: > R\$2500)						
≤ 500	18.02	9.15–35.50	<0.001	2.60	2.34–2.88	<0.001
501 a 1500	10.88	6.23–19.00	<0.001	2.54	2.34–2.76	<0.001
1501 a 2500	4.41	2.39–8.15	<0.001	1.66	1.50–2.20	<0.001
Health insurance (ref.: yes)						
No	8.71	5.77–13.16	<0.001	2.57	2.40–2.75	<0.001
Enabling organisation						
Registered in FHT (ref.: yes)						
No	0.96	0.79–1.16	0.643	0.79	0.71–0.89	<0.001
Unknown	0.99	0.72–1.38	0.970	0.81	0.69–0.95	0.011
Need perceived						
Perceived dental health (ref.: very good / good)						
Regular	1.46	1.20–1.78	<0.001	1.74	1.63–1.85	<0.001
Poor / very poor	3.52	2.55–4.85	<0.001	2.63	2.31–3.00	<0.001
Eating difficulties due to oral problems (ref.: no)						
Yes	2.24	1.69–2.98	<0.001	1.55	1.41–1.71	<0.001

(Continued)

Table 3. (Continued)

Variables	Non-utilisation			Last dental visit > 12 months		
	OR	95% CI	p-value	OR	95% CI	p-value
<i>Need evaluated</i>						
Missing teeth (ref.: no)						
One or more	0.62	0.48–0.81	<0.001	1.09	1.02–1.16	0.015
All	1.54	1.13–2.09	0.006	6.55	5.56–7.71	<0.001

<https://doi.org/10.1371/journal.pone.0192771.t003>

and limitation due to oral problems were only associated with utilisation of dental services over 12 months ago.

Our findings suggest the role of the broader social determinants as barriers to access to dental services for the population even when individual social vulnerability conditions are considered [24]. On the other hand, individual socioeconomic characteristics, including schooling and income, which are byproducts of the social context where people live were meaningful determinants of both dental services outcomes [52]. Previous studies reported that people with high oral health needs were also concentrated in the cities with poor contextual social indicators [13, 15], which suggests an inverse care law—individuals with greater oral health needs are those who face greater difficulties in accessing services.

Similar to previous studies, adults who reported poor oral health showed greater difficulties in the dental services utilisation [26, 29, 31]. With regards to tooth loss, the relationship with dental services utilisation was ambiguous. The odd of non-utilisation of dental services was higher for individuals with no missing teeth. This inverse finding is probably due to the contact with dental services for tooth extraction. Thus, the difficulties in accessing dental services concerning tooth loss are better expressed in the other outcome, since edentulous people were more likely to have visited a dentist over 12 months ago, as expected.

Recent utilisation of dental services was inversely associated with higher age. The literature suggests the decrease in the regular utilisation of services with increasing age [22, 26]. Although our findings may be partly attributed to a cohort and period effects, there is a need to implement effective actions aiming to improve access to dental care among older adults [53]. The difficulties imposed by aging, the decrease in the perception of need due to edentulism and the low availability of public service might make this situation even worse [8]. Behavioural and occupational factors have been suggested to explain why male adults use less the health services [54]. However, some studies have demonstrated that gender differences occur in the intermediate age groups, but not among the youngest [27, 28].

Comparing with other countries [55, 56], few studies in Brazil have reported the association between race/skin colour and utilisation of dental health services. Populational and telephone surveys revealed that adults with white race/skin colour had increased access to dental services than other ethnic groups [7] and the disadvantage of health services utilisation among black and brown comparing with white race/skin colour [10]. This study showed that individuals with black, brown or indigenous race/skin colour had more difficulties in accessing dental services. These findings remained significant after adjustment for schooling and income, which in turn corroborate for the role of racial inequities in the access and utilisation of health services, as previously suggested [10]. Besides access to goods and services, racism is also related to health inequities [57, 58].

The association of education and income with dental services utilisation found in the study has been reported in the literature. Besides, a clear social gradient was observed between lower education and poorer utilisation of dental services [8, 12, 28, 29]. The variable health insurance

Table 4. Hierarchical multilevel analysis for non-utilisation of dental services, Brazil, State capitals, 2013 (n = 27,015).

Variables	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Model 4 OR (95% CI)
Contextual (city-level)				
Predisposing (greater scores indicate better conditions)	0.62 (0.52–0.76)***	0.74 (0.61–0.90)**	0.80 (0.65–0.98)*	0.77 (0.64–0.94)*
Enabling (greater scores indicate better conditions)	0.84 (0.76–0.93)**	0.84 (0.76–0.94)**	0.84 (0.75–0.94)**	0.84 (0.75–0.94)**
Individual level				
Predisposing demographic				
Age (ref.: 18–21)				
22–34		0.75 (0.56–0.99)*	0.80 (0.58–1.10)	1.02 (0.73–1.42)
35–44		0.47 (0.31–0.71)***	0.56 (0.35–0.90)*	0.85 (0.52–1.39)
45–64		0.43 (0.31–0.62)***	0.53 (0.36–0.79)**	0.86 (0.57–1.29)
65+		0.53 (0.34–0.84)**	0.80 (0.48–1.32)	1.18 (0.73–1.92)
Sex (ref.: female)				
Male		1.71 (1.46–2.01)***	1.95 (1.67–2.28)***	1.90 (1.63–2.22)***
Predisposing social				
Race/skin colour (ref.: white)				
Black		1.89 (1.60–2.23)***	1.69 (1.44–2.00)***	1.69 (1.43–1.99)***
Yellow		0.96 (0.34–2.72)	0.90 (0.34–2.40)	0.87 (0.34–2.26)
Brown		1.49 (1.27–1.76)***	1.36 (1.15–1.60)***	1.35 (1.14–1.59)***
Indigenous		3.79 (1.72–8.35)**	3.51 (1.60–7.67)**	3.78 (1.68–8.51)**
Years of schooling (ref.: >11)				
0–4		22.26 (13.03–38.01)***	9.75 (6.05–15.71)***	11.51 (7.06–18.76)***
5–8		10.10 (6.10–16.71)***	4.84 (3.10–7.56)***	5.97 (3.78–9.41)***
9–11		4.24 (2.49–7.22)***	2.46 (1.50–4.04)***	2.77 (1.69–4.56)***
Social network (ref.: IV, High)				
I (Low)		2.40 (1.26–4.56)**	2.01 (1.06–3.82)*	1.86 (1.00–3.46)
II		1.69 (0.91–3.16)	1.47 (0.79–2.74)	1.44 (0.79–2.62)
III		1.66 (0.89–3.08)	1.51 (0.81–2.82)	1.52 (0.84–2.76)
Enabling financing				
Income (ref.: > R\$2500)				
≤ 500			4.52 (2.20–9.27)***	4.45 (2.20–9.03)***
501 a 1500			2.74 (1.54–4.89)**	2.71 (1.54–4.76)**
1501 a 2500			1.87 (0.98–3.58)	1.92 (1.00–3.68)*
Health insurance (ref.: yes)				
No			3.38 (2.21–5.19)***	3.36 (2.19–5.15)***
Need perceived				
Perceived dental health (ref.: very good / good)				
Regular				1.10 (0.90–1.33)
Poor / very poor				2.02 (1.49–2.73)***
Eating difficulties due to oral problems (ref.: no)				
Yes				1.32 (0.96–1.81)
Need evaluated				
Missing teeth (ref.: no)				
One or more				0.35 (0.26–0.48)***
All				0.47 (0.35–0.63)***
Variance at city level (SE)†	0.091 (0.047)	0.098 (0.051)	0.099 (0.054)	0.095 (0.054)

(Continued)

Table 4. (Continued)

Variables	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Model 4 OR (95% CI)
VPC	0.027	0.029	0.029	0.028

Null model: fixed effect, exponentiated intercept = 0.018 (0.014–0.023); random effect, variance (SE) = 0.320 (0.127), $p = 0.011$, VPC = 0.089

* $P < 0.05$

** $P < 0.01$

*** $P < 0.001$

† obtained through random effects

VPC, variance partition coefficient

<https://doi.org/10.1371/journal.pone.0192771.t004>

was not able to discriminate those with and without specific dental coverage, and may be a more general proxy of socioeconomic status. Since it is expected that not all participants that hold health insurance have also dental services coverage, the conditions that led individuals to have the insurance also induce them to use dental health services more easily and with more regularity.

Individuals with small social network were more likely to have a last dental visit over 12 months ago. Although it is not expected that social networks eliminate the different types of barriers related to dental services utilisation, social ties could provide a more frequent contact among those who use the health services. Social connectedness and participation in social groups provide material and emotional support that favours the adoption of healthy behaviours, including frequency of dental visits [59–61]. The effect of social networks on dental services utilisation, however, was limited compared to other social determinants [59].

Despite the effect of the contextual characteristics that reflect the organisation of the oral health services on the utilisation of dental services, associations with the individual variable relating to the enrolment in FHT were not found. In the bivariate analysis, the latter was only associated with recent utilisation, which did not remain in the multivariate analysis, differently from that observed in another study [7]. As already stated, the development of oral health policies in Brazil led to an increase in access to services [1, 5]. Therefore, the assessed items of service organisation in the State capitals seem to be important to deal with the social barriers that preclude the utilisation of dental services, although they had no effect on the regular use of dental services. The low presence of oral health teams in the FHT of large population municipalities may have contributed to this result at the individual level [62]. An ecological study involving 63 countries of all six WHO regions also found no correlation between total health expenditure and the utilisation of dental services in the last 12 months, although it found association of the outcome with dentist-to-population ratio [24], as also described in a multilevel approach [20].

The present study did not assess whether the dental services outcomes were related to public or private health services since our objective was to evaluate the general barriers of dental service access. Another relevant aspect is that the information available from the Brazilian National Health Survey was related to the last dental visit and not to the type of dental service regularly used. Previous studies comparing private and public services have shown that the latter is more sought by individuals with lower schooling, low income and poor oral health conditions [21, 63, 64]. Better infrastructure of oral health services was also associated with greater utilisation of public health services [64]. Countries with public health systems, even if partial coverage, tend to present lower socioeconomic inequalities related to the utilisation of dental health services [65].

Table 5. Hierarchical multilevel analysis for last dental visit over 12 months ago, Brazil, State capitals, 2013 (n = 26,480).

Variables	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Model 4 OR (95% CI)
Contextual (city-level)				
1 Predisposing (greater scores indicate better conditions)	0.87 (0.77–0.99)*	0.95 (0.89–1.03)	0.99 (0.91–1.07)	0.99 (0.92–1.07)
Individual level				
Predisposing demographic				
Age (ref.: 18–21)				
22–34		1.31 (1.18–1.46)***	1.39 (1.25–1.54)***	1.37 (1.23–1.52)***
35–44		1.32 (1.18–1.48)***	1.50 (1.35–1.68)***	1.46 (1.31–1.64)***
45–64		1.47 (1.29–1.68)***	1.79 (1.56–2.06)***	1.72 (1.48–1.99)***
65+		2.39 (2.06–2.77)***	3.26 (2.75–3.86)***	3.17 (2.67–3.77)***
Sex (ref.: female)				
Male		1.42 (1.34–1.51)***	1.52 (1.43–1.61)***	1.50 (1.41–1.59)***
Predisposing social				
Race/skin colour (ref.: white)				
Black		1.18 (1.08–1.30)**	1.09 (0.99–1.20)	1.07 (0.98–1.18)
Yellow		0.95 (0.72–1.24)	0.90 (0.71–1.15)	0.88 (0.69–1.13)
Brown		1.19 (1.11–1.28)***	1.11 (1.03–1.19)**	1.10 (1.03–1.18)**
Indigenous		0.94 (0.68–1.30)	0.86 (0.62–1.18)	0.83 (0.61–1.14)
Years of schooling (ref.: >11)				
0–4		3.70 (3.26–4.19)***	2.25 (2.00–2.52)***	2.14 (1.91–2.40)***
5–8		2.90 (2.60–3.23)***	1.87 (1.70–2.07)***	1.80 (1.64–1.98)***
9–11		1.71 (1.55–1.88)***	1.27 (1.16–1.38)***	1.24 (1.14–1.36)***
Social network (ref.: IV, High)				
I (Low)		1.83 (1.61–2.07)***	1.60 (1.41–1.81)***	1.59 (1.40–1.81)***
II		1.39 (1.24–1.57)***	1.25 (1.11–1.41)***	1.25 (1.11–1.41)***
III		1.19 (1.05–1.36)**	1.11 (0.98–1.27)	1.11 (0.97–1.27)
Enabling financing				
Income (ref.: > R\$2500)				
≤ 500			1.61 (1.43–1.82)***	1.55 (1.37–1.75)***
501 a 1500			1.51 (1.39–1.64)***	1.49 (1.37–1.62)***
1501 a 2500			1.26 (1.13–1.40)***	1.25 (1.13–1.39)***
Health insurance (ref.: yes)				
No			1.81 (1.70–1.92)***	1.76 (1.65–1.87)***
Enabling organisation				
Registered in FHT (ref.: yes)				
No			1.01 (0.93–1.10)	
Unknown			1.04 (0.90–1.20)	
Need perceived				
Perceived dental health (ref.: very good / good)				
Regular				1.36 (1.27–1.47)***
Poor / very poor				1.76 (1.52–2.05)***
Eating difficulties due to oral problems (ref.: no)				
Yes				0.88 (0.79–0.97)*
Need evaluated				
Missing teeth (ref.: no)				
One or more				0.75 (0.70–0.80)***
All				2.85 (2.44–3.34)***

(Continued)

Table 5. (Continued)

Variables	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Model 4 OR (95% CI)
Variance at city level (SE)†	0.050 (0.020)*	0.069 (0.020)***	0.062 (0.019)**	0.063 (0.018)***
VPC	0.015	0.021	0.018	0.019

Null model: fixed effect, exponentiated intercept = 0.963 (0.863–1.074); random effect, variance (SE) = 0.123 (0.030), $p < 0.001$, VPC = 0.036

* $P < 0.05$

** $P < 0.01$

*** $P < 0.001$

† obtained through random effects

VPC, variance partition coefficient

<https://doi.org/10.1371/journal.pone.0192771.t005>

The cross-sectional nature of this study suggests that any causal inferences must be made with caution [50]. Despite the efforts of standardisation and training, information bias might have occurred due to the large team of interviewers participating in the field work data collection of the survey [44]. On the other hand, the large sample size and its representativeness is a strength of the study. The high number of participants in each city also counterpoises for the low number of groups in the second level (<30) for multilevel logistic regression models [66]. The studied sample included adults from the State capitals. Although this approach allowed the analysis of contextual variables in a multilevel model, the findings might not be generalised to other population groups. In addition, this resulted in a high homogeneity that may explain the low percentage variance explained by the contextual level. Nevertheless, it allowed adequate interpretation and greater applicability of the findings, providing relevant information to support the development of intersectoral public policies that can impact oral health and reduce inequalities in the access and utilisation of dental services.

Conclusions

Interventions tackling contextual and individual characteristics can reduce barriers to dental services utilisation and therefore contribute to the improvement of the oral health status of the population, mainly among those individuals with the most need for dental services in the Brazilian State capitals. Therefore, it is paramount to develop new strategies and to reinforce current actions for the active inclusion of groups with less access to dental services, including elderly people, males, those with black, brown and indigenous race/skin colour, as well as individuals with low schooling, low income and edentulous people. The implementation of evidence-based intersectoral actions along with those specific to the health sector can provide a more rational and effective use of available resources.

Author Contributions

Conceptualization: Fernando José Herkrath, Mario Vianna Vettore, Guilherme Loureiro Werneck.

Data curation: Fernando José Herkrath.

Formal analysis: Fernando José Herkrath, Mario Vianna Vettore, Guilherme Loureiro Werneck.

Investigation: Fernando José Herkrath.

Methodology: Fernando José Herkrath, Mario Vianna Vettore, Guilherme Loureiro Werneck.

Supervision: Mario Vianna Vettore, Guilherme Loureiro Werneck.

Writing – original draft: Fernando José Herkrath.

Writing – review & editing: Fernando José Herkrath, Mario Vianna Vettore, Guilherme Loureiro Werneck.

References

1. Pucca Junior GA, Lucena EH, Cawahisa PT. Financing national policy on oral health in Brazil in the context of the Unified Health System. *Braz Oral Res.* 2010; 24 Suppl 1:26–32.
2. Brasil. Diretrizes da Política Nacional de Saúde Bucal. Brasília: Ministério da Saúde; 2004. Available from: http://189.28.128.100/dab/docs/publicacoes/geral/diretrizes_da_politica_nacional_de_saude_bucal.pdf
3. Paim J, Travassos C, Almeida C, Bahia L, Macinko J. The Brazilian health system: history, advances, and challenges. *Lancet.* 2011; 377(9779):1778–97. [https://doi.org/10.1016/S0140-6736\(11\)60054-8](https://doi.org/10.1016/S0140-6736(11)60054-8) PMID: 21561655
4. Macinko J, Harris MJ. Brazil's family health strategy—delivering community-based primary care in a universal health system. *The New England journal of medicine.* 2015; 372(23):2177–81. <https://doi.org/10.1056/NEJMp1501140> PMID: 26039598
5. Pucca Junior GA, Gabriel M, de Araujo ME, de Almeida FC. Ten Years of a National Oral Health Policy in Brazil: Innovation, Boldness, and Numerous Challenges. *J Dent Res.* 2015; 94(10):1333–7. <https://doi.org/10.1177/0022034515599979> PMID: 26316461
6. Bocolini CS, de Souza Junior PR. Inequities in Healthcare utilization: results of the Brazilian National Health Survey, 2013. *Int J Equity Health.* 2016; 15(1):150. <https://doi.org/10.1186/s12939-016-0444-3> PMID: 27852269
7. Mullachery P, Silver D, Macinko J. Changes in health care inequity in Brazil between 2008 and 2013. *Int J Equity Health.* 2016; 15(1):140. <https://doi.org/10.1186/s12939-016-0431-8> PMID: 27852309
8. Matos DL, Giatti L, Lima-Costa MF. Fatores sócio-demográficos associados ao uso de serviços odontológicos entre idosos brasileiros: um estudo baseado na Pesquisa Nacional por Amostra de Domicílios. *Cad Saude Publica.* 2004; 20(5):1290–7. PMID: 15486672
9. Antunes JLF, Narvai PC. Dental health policies in Brazil and their impact on health inequalities. *Rev Saude Publica.* 2010; 44(2):360–5. PMID: 20339637
10. Peres KG, Peres MA, Boing AF, Bertoldi AD, Bastos JL, Barros AJD. Reduction of social inequalities in utilization of dental care in Brazil from 1998 to 2008. *Rev Saude Publica.* 2012; 46(2):250–8. PMID: 22437856
11. Soares CL. Constructing public oral health policies in Brazil: issues for reflection. *Braz Oral Res.* 2012; 26 Suppl 1:94–102.
12. Ferreira CO, Antunes JL, Andrade FB. Factors associated with the use of dental services by elderly Brazilians. *Rev Saude Publica.* 2013; 47 Suppl 3:90–7.
13. Roncalli AG, Tsakos G, Sheiham A, de Souza GC, Watt RG. Social determinants of dental treatment needs in Brazilian adults. *BMC Public Health.* 2014; 14:1097. <https://doi.org/10.1186/1471-2458-14-1097> PMID: 25339315
14. Viacava F, Bellido JG. Health, access to services and sources of payment, according to household surveys. *Cien Saude Colet.* 2016; 21(2):351–70. <https://doi.org/10.1590/1413-81232015212.19422015> PMID: 26910144
15. Goulart MA, Vettore MV. Is the relative increase in income inequality related to tooth loss in middle-aged adults? *J Public Health Dent.* 2016; 76(1):65–75. <https://doi.org/10.1111/jphd.12113> PMID: 26228934
16. Vettore MV, Aqeeli A. The roles of contextual and individual social determinants of oral health-related quality of life in Brazilian adults. *Qual Life Res.* 2016; 25(4):1029–42. <https://doi.org/10.1007/s11136-015-1118-0> PMID: 26342929
17. Hulka BS, Wheat JR. Patterns of utilization: the patient perspective. *Med Care.* 1985; 23(5):438–60. PMID: 4010347
18. Pavão ALB, Coeli CM, Lopes CS, Faerstein E, Werneck GL, Chor D. Social determinants of the use of health services among public university workers. *Rev Saude Publica.* 2012; 46:98–103. PMID: 22218759
19. Garcia-Subirats I, Vargas Lorenzo I, Mogollon-Perez AS, De Paepe P, da Silva MR, Unger JP, et al. Determinants of the use of different healthcare levels in the General System of Social Security in Health

- in Colombia and the Unified Health System in Brazil. *Gac Sanit.* 2014; 28(6):480–8. <https://doi.org/10.1016/j.gaceta.2014.05.010> PMID: 25048392
20. Lee W, Kim SJ, Albert JM, Nelson S. Community factors predicting dental care utilization among older adults. *J Am Dent Assoc.* 2014; 145(2):150–8. <https://doi.org/10.14219/jada.2013.22> PMID: 24487606
21. Pinto RS, Abreu MHNG, Vargas AMD. Comparing adult users of public and private dental services in the state of Minas Gerais, Brazil. *BMC Oral Health.* 2014; 14:100. <https://doi.org/10.1186/1472-6831-14-100> PMID: 25099268
22. Baldani MH, Brito WH, Lawder JAC, Mendes YBE, Silva FFM, Antunes JLF. Determinantes individuais da utilização de serviços odontológicos por adultos e idosos de baixa renda. *Rev Bras Epidemiol.* 2010; 13:150–62. PMID: 20683563
23. Tchicaya A, Lorentz N. Socioeconomic inequalities in the non-use of dental care in Europe. *Int J Equity Health.* 2014; 13:7. <https://doi.org/10.1186/1475-9276-13-7> PMID: 24476233
24. Bhandari B, Newton JT, Bernabé E. Income inequality, disinvestment in health care and use of dental services. *J Public Health Dent.* 2015; 75(1):58–63. <https://doi.org/10.1111/jphd.12071> PMID: 25176508
25. Matos DL, Lima-Costa MFF, Guerra HL, Marcenes W. Projeto Bambuí: estudo de base populacional dos fatores associados com o uso regular de serviços odontológicos em adultos. *Cad Saude Publica.* 2001; 17:661–8. PMID: 11395802
26. Afonso-Souza G, Nadanovsky P, Chor D, Faerstein E, Werneck GL, Lopes CS. Association between routine visits for dental checkup and self-perceived oral health in an adult population in Rio de Janeiro: the Pro-Saude Study. *Community Dent Oral Epidemiol.* 2007; 35(5):393–400. <https://doi.org/10.1111/j.1600-0528.2006.00343.x> PMID: 17822488
27. Manhães ALD, Costa AJL. Acesso a e utilização de serviços odontológicos no Estado do Rio de Janeiro, Brasil, em 1998: um estudo exploratório a partir da Pesquisa Nacional por Amostra de Domicílios. *Cad Saude Publica.* 2008; 24(1):207–18. PMID: 18209849
28. Camargo MJB, Dumith SC, Barros AJD. Uso regular de serviços odontológicos entre adultos: padrões de utilização e tipos de serviços. *Cad Saude Publica.* 2009; 25(9):1894–906. PMID: 19750377
29. Machado LP, Camargo MJB, Jeronymo JCM, Bastos GAN. Regular use of dental services among adults and older adults in a vulnerable region in Southern Brazil. *Rev Saude Publica.* 2012; 46(3):526–33. PMID: 22635038
30. Miranda CD, Peres MA. Determinantes da utilização de serviços odontológicos entre adultos: um estudo de base populacional em Florianópolis, Santa Catarina, Brasil. *Cad Saude Publica.* 2013; 29(11):2319–32. PMID: 24233046
31. Silva AE, Langlois CO, Feldens CA. Use of dental services and associated factors among elderly in southern Brazil. *Rev Bras Epidemiol.* 2013; 16(4):1005–16. PMID: 24896605
32. Gomes AM, Thomaz EB, Alves MT, Silva AA, Silva RA. Fatores associados ao uso dos serviços de saúde bucal: estudo de base populacional em municípios do Maranhão, Brasil. *Cien Saude Colet.* 2014; 19(2):629–40. PMID: 24863839
33. Travassos C, Viacava F. Acesso e uso de serviços de saúde em idosos residentes em áreas rurais, Brasil, 1998 e 2003. *Cad Saude Publica.* 2007; 23:2490–502. PMID: 17891308
34. Peres MA, Iser BPM, Boing AF, Yokota RTC, Malta DC, Peres KG. Desigualdades no acesso e na utilização de serviços odontológicos no Brasil: análise do Sistema de Vigilância de Fatores de Risco e Proteção para Doenças Crônicas por Inquérito Telefônico (VIGITEL 2009). *Cad Saude Publica.* 2012; 28 Suppl:s90–s100.
35. Cruz GD, Chen Y, Salazar CR, Karloopia R, LeGeros RZ. Determinants of oral health care utilization among diverse groups of immigrants in New York City. *J Am Dent Assoc.* 2010; 141(7):871–8. <https://doi.org/10.14219/jada.archive.2010.0286> PMID: 20592408
36. Petersen PE, Bourgeois D, Ogawa H, Estupinan-Day S, Ndiaye C. The global burden of oral diseases and risks to oral health. *Bulletin of the World Health Organization.* 2005; 83(9):661–9. Epub 2005/10/08. PMID: 16211157
37. Finlayson TL, Gansky SA, Shain SG, Weintraub JA. Dental utilization among Hispanic adults in agricultural worker families in California's Central Valley. *J Public Health Dent.* 2010; 70(4):292–9. <https://doi.org/10.1111/j.1752-7325.2010.00184.x> PMID: 20545826
38. Kengne Talla P, Gagnon MP, Dramaix M, Leveque A. Barriers to dental visits in Belgium: a secondary analysis of the 2004 National Health Interview Survey. *J Public Health Dent.* 2013; 73(1):32–40. <https://doi.org/10.1111/jphd.12003> PMID: 23215822
39. Teusner DN, Brennan DS, Spencer AJ. Dental insurance, attitudes to dental care, and dental visiting. *J Public Health Dent.* 2013; 73(2):103–11. <https://doi.org/10.1111/j.1752-7325.2012.00345.x> PMID: 22747628

40. Okunseri C, Garcia RI, Okunseri E, Visotcky A, Szabo A. Dental service utilization and neighborhood characteristics in young adults in the United States: a multilevel approach. *J Public Health Dent*. 2015; 75(4):282–90. <https://doi.org/10.1111/jphd.12097> PMID: [25929275](#)
41. Bhandari B, Newton JT, Bernabe E. Income Inequality and Use of Dental Services in 66 Countries. *J Dent Res*. 2015; 94(8):1048–54. <https://doi.org/10.1177/0022034515586960> PMID: [25994178](#)
42. Kassebaum NJ, Smith AGC, Bernabe E, Fleming TD, Reynolds AE, Vos T, et al. Global, Regional, and National Prevalence, Incidence, and Disability-Adjusted Life Years for Oral Conditions for 195 Countries, 1990–2015: A Systematic Analysis for the Global Burden of Diseases, Injuries, and Risk Factors. *Journal of dental research*. 2017; 96(4):380–7. Epub 2017/08/10. <https://doi.org/10.1177/0022034517693566> PMID: [28792274](#)
43. Andersen RM, Davidson PL. Improving access to care in America: individual and contextual indicators. In: Kominski GF, editor. *Changing the US Health Care System: Key Issues in Health Services Policy and Management* 4th ed. San Francisco: Jossey-Bass; 2014. p. 3–31.
44. Szwarcwald CL, Malta DC, Pereira CA, Vieira ML, Conde WL, Souza Junior PR, et al. Pesquisa Nacional de Saúde no Brasil: concepção e metodologia de aplicação. *Cien Saude Colet*. 2014; 19(2):333–42. PMID: [24863810](#)
45. Travassos C, Williams DR. The concept and measurement of race and their relationship to public health: a review focused on Brazil and the United States. *Cad Saude Publica*. 2004; 20(3):660–78. PMID: [15263977](#)
46. Berkman LF, Syme SL. Social networks, host resistance, and mortality: a nine-year follow-up study of Alameda County residents. *Am J Epidemiol*. 1979; 109(2):186–204. PMID: [425958](#)
47. United Nations Development Programme. Human Development Atlas in Brazil 2013 [cited 2017]. Available from: <http://www.atlasbrasil.org.br/2013/en/home/>.
48. Ministério da Saúde. Datasus—Departamento de Informática do SUS [cited 2017]. Available from: <http://datasus.saude.gov.br/>.
49. Ministério da Saúde. Secretaria de Atenção à Saúde. Secretaria de Vigilância em Saúde. Departamento de Atenção Básica. Coordenação Nacional de Saúde Bucal. SBBrazil 2010: Pesquisa Nacional de Saúde Bucal: Ministério da Saúde; 2012 [cited 2017]. Available from: http://bvsms.saude.gov.br/bvs/publicacoes/pesquisa_nacional_saude_bucal.pdf.
50. Reichenheim ME, Coutinho ES. Measures and models for causal inference in cross-sectional studies: arguments for the appropriateness of the prevalence odds ratio and related logistic regression. *BMC Med Res Methodol*. 2010; 10:66. <https://doi.org/10.1186/1471-2288-10-66> PMID: [20633293](#)
51. Snijders TAB, Bosker RJ. Multilevel analysis: An introduction to basic and advanced multilevel modeling (2nd ed.). Los Angeles/London/New Delhi/Singapore/Washington D.C.: SAGE Publications Inc.; 2012.
52. Andrew MK, Keefe JM. Social vulnerability from a social ecology perspective: a cohort study of older adults from the National Population Health Survey of Canada. *BMC Geriatr*. 2014; 14:90. <https://doi.org/10.1186/1471-2318-14-90> PMID: [25129548](#)
53. Astrom AN, Ekback G, Nasir E, Ordell S, Unell L. Use of dental services throughout middle and early old ages: a prospective cohort study. *Community Dent Oral Epidemiol*. 2013; 41(1):30–9. <https://doi.org/10.1111/j.1600-0528.2012.00709.x> PMID: [22937755](#)
54. Muirhead VE, Quinonez C, Figueiredo R, Locker D. Predictors of dental care utilization among working poor Canadians. *Community Dent Oral Epidemiol*. 2009; 37(3):199–208. <https://doi.org/10.1111/j.1600-0528.2009.00471.x> PMID: [19508268](#)
55. Flores G, Lin H. Trends in racial/ethnic disparities in medical and oral health, access to care, and use of services in US children: has anything changed over the years? *Int J Equity Health*. 2013; 12:10. <https://doi.org/10.1186/1475-9276-12-10> PMID: [23339566](#)
56. Horner-Johnson W, Dobberty K, Beilstein-Wedel E. Disparities in dental care associated with disability and race and ethnicity. *J Am Dent Assoc*. 2015; 146(6):366–74. <https://doi.org/10.1016/j.adaj.2015.01.024> PMID: [26025823](#)
57. Gee GC, Ford CL. STRUCTURAL RACISM AND HEALTH INEQUITIES: Old Issues, New Directions. *Du Bois Rev*. 2011; 8(1):115–32. Epub 2011/04/01. <https://doi.org/10.1017/S1742058X11000130> PMID: [25632292](#)
58. Williams DR. Miles to go before we sleep: racial inequities in health. *J Health Soc Behav*. 2012; 53(3):279–95. <https://doi.org/10.1177/0022146512455804> PMID: [22940811](#)
59. Sabbah W, Tsakos G, Chandola T, Newton T, Kawachi I, Sheiham A, et al. The relationship between social network, social support and periodontal disease among older Americans. *J Clin Periodontol*. 2011; 38(6):547–52. <https://doi.org/10.1111/j.1600-051X.2011.01713.x> PMID: [21362014](#)

60. Watt RG, Heilmann A, Sabbah W, Newton T, Chandola T, Aida J, et al. Social relationships and health related behaviors among older US adults. *BMC Public Health*. 2014; 14:533. <https://doi.org/10.1186/1471-2458-14-533> PMID: [24885507](https://pubmed.ncbi.nlm.nih.gov/24885507/)
61. Gulcan F, Ekback G, Ordell S, Lie SA, Astrom AN. Social predictors of less frequent dental attendance over time among older people: population-averaged and person-specific estimates. *Community Dent Oral Epidemiol*. 2016; 44(3):263–73. <https://doi.org/10.1111/cdoe.12214> PMID: [26854281](https://pubmed.ncbi.nlm.nih.gov/26854281/)
62. Lucena EHG, Pucca Júnior GA, Sousa MF. A política nacional de saúde bucal no Brasil no contexto do Sistema Único de Saúde. *Tempus: Actas de Saúde Coletiva*. 2011; 5(3):53–63.
63. Pinto RS, Matos DL, Loyola Filho AI. Características associadas ao uso de serviços odontológicos públicos pela população adulta brasileira. *Cien Saude Colet*. 2012; 17(2):531–44. PMID: [22267047](https://pubmed.ncbi.nlm.nih.gov/22267047/)
64. Pinto RS, Roncalli AG, Abreu MH, Vargas AM. Use of Public Oral Health Services by the Adult Population: A Multilevel Analysis. *PLoS One*. 2016; 11(1):e0145149. <https://doi.org/10.1371/journal.pone.0145149> PMID: [26730714](https://pubmed.ncbi.nlm.nih.gov/26730714/)
65. Palencia L, Espelt A, Cornejo-Ovalle M, Borrell C. Socioeconomic inequalities in the use of dental care services in Europe: what is the role of public coverage? *Community Dent Oral Epidemiol*. 2014; 42(2):97–105. <https://doi.org/10.1111/cdoe.12056> PMID: [23786417](https://pubmed.ncbi.nlm.nih.gov/23786417/)
66. Moineddin R, Matheson FI, Glazier RH. A simulation study of sample size for multilevel logistic regression models. *BMC Med Res Methodol*. 2007; 7:34. <https://doi.org/10.1186/1471-2288-7-34> PMID: [17634107](https://pubmed.ncbi.nlm.nih.gov/17634107/)